

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.


Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

A



B






Figure.1: A- Nestin positive cells;

B- Expansion of Nestin positive cells



Histogram Statistics
 File: NEURAL(30/9/02) 001
 Sample ID: CONTROL/NCAM
 Acquisition Date: 30-Sep-02 Gate: G1
 Gated Events: 30448 Total Events: 50000
 Marker Events % Gated % Total
 All 30448 10000 60.90
 M1 165 054 033



Histogram Statistics
 File: NEURAL(30/9/02) 004
 Sample ID: NCAM(FIXED/PERM)
 Acquisition Date: 30-Sep-02 Gate: G1
 Gated Events: 32566 Total Events: 50000
 Marker Events % Gated % Total
 All 32566 10000 65.13
 M1 31434 9652 62.87

Figure.2: FACS analysis


A-Control (unlabelled cells)

B- NCAM positive cells

A



B



**Figure. 3 : Magnetic Sorting of NCAM positive cells
A, B- NCAM positive cells**




Figure.4A : MAP-2 positive cells





Figure.4B : β Tubulin positive cells

A



B




Figure.5 : Cells positive for Tyrosine hydroxylase




Figure.6 Oligodendrocyte






Figure.7 Serotonin




TH / β tubulin



TH/MAP-2



TH/Nurrl



TH /DAT

Figure. 8. Co-localisation of TH with other neuronal antigens




Figure 9:Percent proportion of TH positive neurons





Figure 10 : Quantitative analysis of neuronal populations from NCAM enriched cells.
Tyrosine hydroxylase positive cells found to be 60%



**Figure 11 : Quantitative analysis of neuronal populations from Nestin expansion and differentiation.
Tyrosine hydroxylase positive cells found to be 40%**




Figure. 12 : Expression of Dopamine specific genes in ES derived neurons




Figure. 13 : Analysis of intracellular dopamine in cell lysates




Figure14: KCl evoked release of dopamine in conditioned medium